

FACT: One out of every 4 children attending school has been exposed to a traumatic event that can affect learning and/or behavior.

FACT: Trauma can impact school performance.

- Lower GPA
- Higher rate of school absences
- Increased drop-out
- More suspensions and expulsions
- Decreased reading ability

FACT: Trauma can impair learning.

Single exposure to traumatic events may cause jumpiness, intrusive thoughts, interrupted sleep and nightmares, anger and moodiness, and/or social withdrawal—any of which can interfere with concentration and memory.

Chronic exposure to traumatic events, especially during a child’s early years, can:

- Adversely affect attention, memory, and cognition
- Reduce a child’s ability to focus, organize, and process information
- Interfere with effective problem solving and/or planning
- Result in overwhelming feelings of frustration and anxiety

FACT: Traumatized children may experience physical and emotional distress.

- Physical symptoms like headaches and stomachaches
- Poor control of emotions
- Inconsistent academic performance
- Unpredictable and/or impulsive behavior
- Over or under-reacting to bells, physical contact, doors slamming, sirens, lighting, sudden movements
- Intense reactions to reminders of their traumatic event:
 - Thinking others are violating their personal space, i.e., “What are you looking at?”
 - Blowing up when being corrected or told what to do by an authority figure
 - Fighting when criticized or teased by others
 - Resisting transition and/or change

FACT: You can help a child who has been traumatized.

- Follow your school’s reporting procedures if you suspect abuse
- Work with the child’s caregiver(s) to share and address school problems
- Refer to community resources when a child shows signs of being unable to cope with traumatic stress
- Share Trauma Facts for Educators with other teachers and school personnel

This project was funded by the Substance Abuse and Mental Health Services Administration (SAMHSA), US Department of Health and Human Services (HHS). The views, policies, and opinions expressed are those of the authors and do not necessarily reflect those of SAMHSA or HHS.